天成醫療體系

化療藥品衛教單張（醫護版）
愛斯達好克癌注射劑（**Holoxan**）

外觀	商品名	中文名	學名	規格含量
	Holoxan	愛斯達好克癌注射劑	Ifosfamide	2g/vial

給藥方式：IVF
懷孕分級：D（FDA）、D（AUS）
授乳分級：Infant risk cannot be ruled out.

一、作用機轉：
Ifosfamide is classified as an alkylating agent of the nitrogen mustard type. After metabolic activation, active metabolites of ifosfamide alkylate or bind with many intracellular molecular structures, including nucleic acids. The cytotoxic action is primarily due to cross-linking of strands of DNA and RNA, as well as inhibition of protein synthesis

二、原液保存條件：室溫避光存放 。

三、配製及保存期限：以 50ml water for injection 配製

四、稀釋及保存期限：以D_5W稀釋，配製好的溶液如放在溫度不超過8℃的冰箱內，則可保存24小時。

五、其他注意事項：
1. 腎功能不全者減量；增加液體的攝取，有助於避免出血性膀胱炎。
2. 當使用高劑量治療時，Mesna 對預防該併發症也有益。
3. 以緩慢間歇性靜脈點滴給藥至少需要 30 分鐘以上且給藥最終濃度不可超過 40mg/ml(通常給藥濃度是在 0.6-20mg/ml 之間)，或可以 24 小時的點滴給藥。

六、外滲處理方式：局部冷敷:每次敷 60 分鐘,8 小時一次,共 3 天。